


Atoms, elements and compounds

Atom	<i>The smallest part of an element that can exist</i>	Have a radius of around 0.1 nanometres and have no charge (0).
Element	<i>Contains only one type of atom</i>	Around 100 different elements each one is represented by a symbol e.g. O, Na, Br.
Compound	<i>Two or more elements chemically combined</i>	Compounds can only be separated into elements by chemical reactions.



Central nucleus	Contains protons and neutrons
Electron shells	Contains electrons

Name of Particle	Relative Charge	Relative Mass
Proton	+1	1
Neutron	0	1
Electron	-1	Very small

Electronic shell	Max number of electrons
1	2
2	8
3	8
4	2

Electronic structures

AQA Trilogy
C1: Atomic Structure & The Periodic Table

Relative electrical charges of subatomic particles

7 Li	Mass number	<i>The sum of the protons and neutrons in the nucleus</i>	
3	Atomic number	<i>The number of protons in the atom</i>	Number of electrons = number of protons

Mixtures	<i>Two or more elements or compounds not chemically combined together</i>	Can be separated by physical processes.
----------	---	---

Method	Description	Example
Filtration	<i>Separating an insoluble solid from a liquid</i>	To get sand from a mixture of sand, salt and water.
Crystallisation	<i>To separate a solid from a solution</i>	To obtain pure crystals of sodium chloride from salt water.
Simple distillation	<i>To separate a solvent from a solution</i>	To get pure water from salt water.
Fractional distillation	<i>Separating a mixture of liquids each with different boiling points</i>	To separate the different compounds in crude oil.
Chromatography	<i>Separating substances that move at different rates through a medium</i>	To separate out the dyes in food colouring.

Pre 1900		<i>Tiny solid spheres that could not be divided</i>	Before the discovery of the electron, John Dalton said the solid sphere made up the different elements.
1897 'plum pudding'		<i>A ball of positive charge with negative electrons embedded in it</i>	JJ Thompson 's experiments showed that showed that an atom must contain small negative charges (discovery of electrons).
1909 nuclear model		<i>Positively charge nucleus at the centre surrounded negative electrons</i>	Ernest Rutherford's alpha particle scattering experiment showed that the mass was concentrated at the centre of the atom.
1913 Bohr model		<i>Electrons orbit the nucleus at specific distances</i>	Niels Bohr proposed that electrons orbited in fixed shells; this was supported by experimental observations.

The development of the model of the atom	James Chadwick	<i>Provided the evidence to show the existence of neutrons within the nucleus</i>
--	----------------	---

Rutherford's scattering experiment	<i>A beam of alpha particles are directed at a very thin gold foil</i>		Most of the alpha particles passed right through. A few (+) alpha particles were deflected by the positive nucleus. A tiny number of particles reflected back from the nucleus.

Chemical equations	<i>Show chemical reactions - need reactant(s) and product(s) energy always involves and energy change</i>	Law of conservation of mass states the total mass of products = the total mass of reactants.
--------------------	---	--

Word equations	<i>Uses words to show reaction</i> reactants → products magnesium + oxygen → magnesium oxide	Does not show what is happening to the atoms or the number of atoms.
Symbol equations	<i>Uses symbols to show reaction</i> reactants → products 2Mg + O ₂ → 2MgO	Shows the number of atoms and molecules in the reaction, these need to be balanced.

Relative atomic mass	Isotopes	<i>Atoms of the same element with the same number of protons and different numbers of neutrons</i>	³⁵Cl (75%) and ³⁷Cl (25%) Relative abundance = (% isotope 1 x mass isotope 1) + (% isotope 2 x mass isotope 2) ÷ 100 e.g. (25 x 37) + (75x 35) ÷ 100 = 35.5



Alkali metals: 1, 2
 Halogens: 3, 4, 5, 6, 7
 Noble gases: 0

H	Transition metals																He						
Li	Be																	B	C	N	O	F	Ne
Na	Mg																	Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr						
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe						
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn						
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	?	?	?												

Elements arranged in order of atomic number

Elements with similar properties are in columns called groups

Elements in the same group have the same number of outer shell electrons and elements in the same period (row) have the same number of electron shells.

The Periodic table

Development of the Periodic table

Before discovery of protons, neutrons and electrons	Elements arranged in order of atomic weight	Early periodic tables were incomplete, some elements were placed in inappropriate groups if the strict order of atomic weights was followed.
Mendeleev	Left gaps for elements that hadn't been discovered yet	Elements with properties predicted by Mendeleev were discovered and filled in the gaps. Knowledge of isotopes explained why order based on atomic weights was not always correct.

Metals to the left of this line, non metals to the right

Metals	To the left of the Periodic table	Form positive ions. Conductors, high melting and boiling points, ductile, malleable.
Non metals	To the right of the Periodic table	Form negative ions. Insulators, low melting and boiling points.

Metals and non metals

AQA Trilogy
 C1: Atomic Structure & The Periodic Table

Group 7

Group 1

Alkali metals	Very reactive with oxygen, water and chlorine	Only have one electron in their outer shell. Form +1 ions.
	Reactivity increases down the group	Negative outer electron is further away from the positive nucleus so is more easily lost.

Halogens	Consist of molecules made of a pair of atoms	Have seven electrons in their outer shell. Form -1 ions.
	Melting and boiling points increase down the group (gas → liquid → solid)	Increasing atomic mass number.
	Reactivity decreases down the group	Increasing proton number means an electron is more easily gained

Group 0

Noble gases	Unreactive, do not form molecules	This is due to having full outer shells of electrons.
	Boiling points increase down the group	Increasing atomic number.

With metals	Forms a metal halide	Metal + halogen → metal halide e.g. Sodium + chlorine → sodium chloride	e.g. NaCl metal atom loses outer shell electrons and halogen gains an outer shell electron
With hydrogen	Forms a hydrogen halide	Hydrogen + halogen → hydrogen halide e.g. Hydrogen + bromine → hydrogen bromide	e.g. Cl ₂ + H ₂ → 2HCl
With aqueous solution of a halide salt	A more reactive halogen will displace the less reactive halogen from the salt	Chlorine + potassium bromide → potassium chloride + bromine	e.g. Cl ₂ + 2KBr → 2KCl + Br ₂

With oxygen	Forms a metal oxide	Metal + oxygen → metal oxide	e.g. 4Na + O ₂ → 2Na ₂ O
With water	Forms a metal hydroxide and hydrogen	Metal + water → metal hydroxide + hydrogen	e.g. 2Na + 2H ₂ O → 2NaOH + H ₂
With chlorine	Forms a metal chloride	Metal + chlorine → metal chloride	e.g. 2Na + Cl ₂ → 2NaCl

Ionic	Particles are oppositely charged ions	Occurs in compounds formed from metals combined with non metals.
Covalent	Particles are atoms that share pairs of electrons	Occurs in most non metallic elements and in compounds of non metals.
Metallic	Particles are atoms which share delocalised electrons	Occurs in metallic elements and alloys.

Solid, liquid, gas

Melting and freezing happen at melting point, boiling and condensing happen at boiling point.

The amount of energy needed for a state change depends on the strength of forces between particles in the substance.

(HT only) Limitations of simple model:

- There are no forces in the model
- All particles are shown as spheres
 - Spheres are solid

s	solid
l	liquid
g	gas

<i>High melting and boiling points</i>	Large amounts of energy needed to break the bonds.
<i>Do not conduct electricity when solid</i>	Ions are held in a fixed position in the lattice and cannot move.
<i>Do conduct electricity when molten or dissolved</i>	Lattice breaks apart and the ions are free to move.

Properties of ionic compounds

Chemical bonds

The three states of matter

<i>Good conductors of electricity</i>	Delocalised electrons carry electrical charge through the metal.
<i>Good conductors of thermal energy</i>	Energy is transferred by the delocalised electrons.

<i>High melting and boiling points</i>	This is due to the strong metallic bonds.
<i>Pure metals can be bent and shaped</i>	Atoms are arranged in layers that can slide over each other.

AQA Trilogy C2: Bonding, Structure & The Properties of Matter

Metals as conductors

Properties of metals and alloys

Alloys	<i>Mixture of two or more elements at least one of which is a metal</i>	Harder than pure metals because atoms of different sizes disrupt the layers so they cannot slide over each other.
---------------	---	---

Pure metal **Alloy**

Metallic bonding

Ionic bonding

Electrons are transferred so that all atoms have a noble gas configuration (full outer shells).	<i>Metal atoms lose electrons and become positively charged ions</i>	Group 1 metals form +1 ions Group 2 metals form +2 ions
	<i>Non metals atoms gain electrons to become negatively charged ions</i>	Group 6 non metals form -2 ions Group 7 non metals form -1 ions

Ionic compounds

Dot and cross diagram

$(2, 8, 1)$ $(2, 8, 7)$ $(2, 8)$ $(2, 8, 8)$

Giant structure

● Na⁺ ● Cl⁻

Structure

- Held together by strong electrostatic forces of attraction between oppositely charged ions
- Forces act in all directions in the lattice

Giant structure of atoms arranged in a regular pattern

Delocalised electrons Metal ions

Electrons in the outer shell of metal atoms are delocalised and free to move through the whole structure. This sharing of electrons leads to strong metallic bonds.



M_r	<i>The sum of the relative atomic masses of the atoms in the numbers shown in the formula</i>	The sum of the M_r of the reactants in the quantities shown equals the sum of the M_r of the products in the quantities shown.	$2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$ $\downarrow \quad \downarrow \quad \downarrow$ $48\text{g} + 32\text{g} = 80\text{g}$ $80\text{g} = 80\text{g}$	The reactant that is completely used up	<i>Limits the amount of product that is made</i>	Less moles of product are made.
-------------------------	---	--	---	---	--	---------------------------------

Mass appears to increase during a reaction	<i>One of the reactants is a gas</i>	Magnesium + oxygen \rightarrow magnesium oxide
Mass appears to decrease during a reaction	<i>One of the products is a gas and has escaped</i>	Calcium carbonate \rightarrow carbon dioxide + calcium oxide

Mass changes when a reactant or product is a gas

Conservation of mass	<i>No atoms are lost or made during a chemical reaction</i>	Mass of the products equals the mass of the reactants.
----------------------	---	--

Balanced symbol equations	<i>Represent chemical reactions and have the same number of atoms of each element on both sides of the equation</i>	$\text{H}_2 + \text{Cl}_2 \rightarrow 2\text{HCl}$ <p>Subscript \rightarrow Normal script</p>
		<p>Subscript numbers show the number of atoms of the element to its left.</p> <p>Normal script numbers show the number of molecules.</p>

Chemical amounts are measured in moles (mol)	<i>Mass of one mole of a substance in grams = relative formula mass</i>	<p>One mole of $\text{H}_2\text{O} = 18\text{g} (1 + 1 + 16)$</p> <p>One mole of $\text{Mg} = 24\text{g}$</p>
--	---	---

Avogadro constant	<i>One mole of any substance will contain the same number of particles, atoms, molecules or ions.</i>	<p>6.02×10^{23} per mole</p> <p>One mole of H_2O will contain 6.02×10^{23} molecules</p> <p>One mole of NaCl will contain 6.02×10^{23} Na^+ ions</p>
-------------------	---	---

<i>Number of moles = $\frac{\text{mass (g)}}{A_r}$ or $\frac{\text{mass (g)}}{M_r}$</i>	<p>How many moles of sulfuric acid molecules are there in 4.7g of sulfuric acid (H_2SO_4)? Give your answer to 1 significant figure.</p> $\frac{4.7}{98} = 0.05 \text{ mol}$ <p style="text-align: center;">\leftarrow (M_r of H_2SO_4)</p>
---	---

AQA Trilogy
C3: Quantitative Chemistry

Relative formula mass (M_r)

Limiting reactants (HT only)

Chemical measurements

Whenever a measurement is taken, there is always some uncertainty about the result obtained	<i>Can determine whether the mean value falls within the range of uncertainty of the result</i>	<ol style="list-style-type: none"> Calculate the mean Calculate the range of the results Estimate of uncertainty in mean would be half the range
---	---	---

Concentration of solutions	<p>Example:</p> <ol style="list-style-type: none"> Mean value is 46.5s Range of results is 44s to 49s = 5s Time taken was $46.5\text{s} \pm 2.5\text{s}$
----------------------------	--

Measured in mass per given volume of solution (g/dm^3)	<i>Conc. = $\frac{\text{mass (g)}}{\text{volume (dm}^3\text{)}}$</i>	<p>HT only</p> <p>Greater mass = higher concentration.</p> <p>Greater volume = lower concentration.</p>
--	---	---

The balancing numbers in a symbol equation can be calculated from the masses of reactants and products	<i>Convert the masses in grams to amounts in moles and convert the number of moles to simple whole number ratios.</i>
--	---

Chemical equations show the number of moles reacting and the number of moles made	<p>$\text{Mg} + 2\text{HCl} \rightarrow \text{MgCl}_2 + \text{H}_2$</p> <p><i>One mole of magnesium reacts with two moles of hydrochloric acid to make one mole of magnesium chloride and one mole of hydrogen</i></p> <p>If you have a 60g of Mg, what mass of HCl do you need to convert it to MgCl_2?</p> <p>A_r : Mg = 24 so mass of 1 mole of Mg = 24g</p> <p>M_r : HCl (1 + 35.5) so mass of 1 mole of HCl = 36.5g</p> <p>So 60g of Mg is $60/24 = 2.5$ moles</p> <p>Balanced symbol equation tells us that for every one mole of Mg, you need two moles of HCl to react with it.</p> <p>So you need $2.5 \times 2 = 5$ moles of HCl</p> <p>You will need $5 \times 36.5\text{g}$ of HCl = 182.5g</p>
---	---



<p>A measure of the amount of starting materials that end up as useful products</p>	<p>Atom economy = $\frac{\text{Relative formula mass of desired product from equation}}{\text{Sum of relative formula mass of all reactants from equation}} \times 100$</p>	<p>High atom economy is important or sustainable development and economic reasons</p>
---	--	---

Calculate the atom economy for making hydrogen by reacting zinc with hydrochloric acid:

$$\text{Zn} + 2\text{HCl} \rightarrow \text{ZnCl}_2 + \text{H}_2$$

M_r of $\text{H}_2 = 1 + 1 = 2$
 M_r of $\text{Zn} + 2\text{HCl} = 65 + 1 + 1 + 35.5 + 35.5 = 138$

Atom economy = $\frac{2}{138} \times 100$
 = $\frac{2}{138} \times 100 = 1.45\%$

This method is unlikely to be chosen as it has a low atom economy.

Atom economy

AQA Trilogy
C3: Quantitative Chemistry

HT only:
 200g of calcium carbonate is heated. It decomposes to make calcium oxide and carbon dioxide. Calculate the theoretical mass of calcium oxide made.

$$\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$$

M_r of $\text{CaCO}_3 = 40 + 12 + (16 \times 3) = 100$
 M_r of $\text{CaO} = 40 + 16 = 56$
 100g of CaCO_3 would make 56 g of CaO
 So 200g would make 112g

Percentage yield

<p>Yield is the amount of product obtained</p>	<p><i>It is not always possible to obtain the calculated amount of a product</i></p>	<p>The reaction may not go to completion because it is reversible.</p> <p>Some of the product may be lost when it is separated from the reaction mixture.</p> <p>Some of the reactants may react in ways different to the expected reaction.</p>
--	--	--

<p>Percentage yield is comparing the amount of product obtained as a percentage of the maximum theoretical amount</p>	<p>$\% \text{ Yield} = \frac{\text{Mass of product made} \times 100}{\text{Max. theoretical mass}}$</p>	<p>A piece of sodium metal is heated in chlorine gas. A maximum theoretical mass of 10g for sodium chloride was calculated, but the actual yield was only 8g.</p> <p>Calculate the percentage yield.</p> <p>Percentage yield = $8/10 \times 100 = 80\%$</p>
---	--	--



Oxidation Is Loss (of electrons) Reduction Is Gain (of electrons)

C4

Ionic half equations (HT only)

For displacement reactions

Ionic half equations show what happens to each of the reactants during reactions

For example:
The ionic equation for the reaction between iron and copper (II) ions is:
 $Fe + Cu^{2+} \rightarrow Fe^{2+} + Cu$

The half-equation for iron (II) is:
 $Fe \rightarrow Fe^{2+} + 2e^{-}$

The half-equation for copper (II) ions is:
 $Cu^{2+} + 2e^{-} \rightarrow Cu$

HT ONLY: Reactions between metals and acids are redox reactions as the metal donates electrons to the hydrogen ions. This displaces hydrogen as a gas while the metal ions are left in the solution.

Reactions with acids	$metal + acid \rightarrow metal\ salt + hydrogen$	magnesium + hydrochloric acid \rightarrow magnesium chloride + hydrogen zinc + sulfuric acid \rightarrow zinc sulfate + hydrogen
-----------------------------	---	---

Acids react with some metals to produce salts and hydrogen.

Reactions of acids and metals

Extraction using carbon

Metals less reactive than carbon can be extracted from their oxides by reduction.

For example:
zinc oxide + carbon \rightarrow zinc + carbon dioxide

Unreactive metals, such as gold, are found in the Earth as the metal itself. They can be mined from the ground.

Reactions of acids

Extraction of metals and reduction

AQA Trilogy C4: Chemical Changes

Reactivity of metals

	Reactions with water	Reactions with acid
Group 1 metals	<i>Reactions get more vigorous as you go down the group</i>	<i>Reactions get more vigorous as you go down the group</i>
Group 2 metals	<i>Do not react with water</i>	<i>Observable reactions include fizzing and temperature increases</i>
Zinc, iron and copper	<i>Do not react with water</i>	<i>Zinc and iron react slowly with acid. Copper does not react with acid.</i>

Oxidation and reduction in terms of electrons (HT ONLY)

Neutralisation of acids and salt production

Acid name	Salt name
Hydrochloric acid	Chloride
Sulfuric acid	Sulfate
Nitric acid	Nitrate

sodium hydroxide + hydrochloric acid \rightarrow sodium chloride + water
calcium carbonate + sulfuric acid \rightarrow calcium sulfate, + carbon dioxide + water

The reactivity series

Metal oxides

Neutralisation

Acids can be neutralised by alkalis and bases

An alkali is a soluble base e.g. metal hydroxide.
A base is a substance that neutralises an acid e.g. a soluble metal hydroxide or a metal oxide.

Metals form positive ions when they react	<i>The reactivity of a metal is related to its tendency to form positive ions</i>	The reactivity series arranges metals in order of their reactivity (their tendency to form positive ions).
Carbon and hydrogen	<i>Carbon and hydrogen are non-metals but are included in the reactivity series</i>	These two non-metals are included in the reactivity series as they can be used to extract some metals from their ores, depending on their reactivity.
Displacement	<i>A more reactive metal can displace a less reactive metal from a compound.</i>	Silver nitrate + Sodium chloride \rightarrow Sodium nitrate + Silver chloride

- potassium most reactive K
sodium Na
calcium Ca
magnesium Mg
aluminium Al
carbon C
zinc Zn
iron Fe
tin Sn
lead Pb
hydrogen H
copper Cu
silver Ag
gold Au
platinum least reactive Pt
-

Metals and oxygen	<i>Metals react with oxygen to form metal oxides</i>	magnesium + oxygen \rightarrow magnesium oxide $2Mg + O_2 \rightarrow 2MgO$
Reduction	<i>This is when oxygen is removed from a compound during a reaction</i>	e.g. metal oxides reacting with hydrogen, extracting low reactivity metals
Oxidation	<i>This is when oxygen is gained by a compound during a reaction</i>	e.g. metals reacting with oxygen, rusting of iron



The ions discharged when an aqueous solution is electrolysed using inert electrodes depend on the relative reactivity of the elements involved.

At the negative electrode	Metal will be produced on the electrode if it is less reactive than hydrogen. Hydrogen will be produced if the metal is more reactive than hydrogen.
At the positive electrode	Oxygen is formed at positive electrode. If you have a halide ion (Cl ⁻ , I ⁻ , Br ⁻) then you will get chlorine, bromine or iodine formed at that electrode.


Process of electrolysis	<i>Splitting up using electricity</i>	When an ionic compound is melted or dissolved in water, the ions are free to move. These are then able to conduct electricity and are called electrolytes. Passing an electric current through electrolytes causes the ions to move to the electrodes.
Electrode	<i>Anode Cathode</i>	The positive electrode is called the anode. The negative electrode is called the cathode.
Where do the ions go?	<i>Cations Anions</i>	Cations are positive ions and they move to the negative cathode. Anions are negative ions and they move to the positive anode.

Extracting metals using electrolysis	<i>Metals can be extracted from molten compounds using electrolysis.</i>
	<i>This process is used when the metal is too reactive to be extracted by reduction with carbon.</i>
	<i>The process is expensive due to large amounts of energy needed to produce the electrical current. Example: aluminium is extracted in this way.</i>

Electrolysis of aqueous solutions

Strong acids	<i>Completely ionised in aqueous solutions e.g. hydrochloric, nitric and sulfuric acids.</i>
Weak acids	<i>Only partially ionised in aqueous solutions e.g. ethanoic acid, citric acid.</i>
Hydrogen ion concentration	<i>As the pH decreases by one unit (becoming a stronger acid), the hydrogen ion concentration increases by a factor of 10.</i>




Soluble salts	<i>Soluble salts can be made from reacting acids with solid insoluble substances (e.g. metals, metal oxides, hydroxides and carbonates).</i>
Production of soluble salts	<i>Add the solid to the acid until no more dissolves. Filter off excess solid and then crystallise to produce solid salts.</i>



You can use universal indicator or a pH probe to measure the acidity or alkalinity of a solution against the pH scale.

In neutralisation reactions, hydrogen ions react with hydroxide ions to produce water:
 $H^+ + OH^- \rightarrow H_2O$

Acids	<i>Acids produce hydrogen ions (H⁺) in aqueous solutions.</i>
Alkalis	<i>Aqueous solutions of alkalis contain hydroxide ions (OH⁻).</i>



Higher tier: You can display what is happening at each electrode using half-equations:
At the cathode: $Pb^{2+} + 2e^- \rightarrow Pb$
At the anode: $2Br^- \rightarrow Br_2 + 2e^-$



Endothermic	Energy is taken in from the surroundings so the temperature of the surroundings decreases	<ul style="list-style-type: none"> Thermal decomposition Sports injury packs
Exothermic	Energy is transferred to the surroundings so the temperature of the surroundings increases	<ul style="list-style-type: none"> Combustion Hand warmers Neutralisation

Reaction profiles	Show the overall energy change of a reaction
-------------------	--

Types of reaction

Breaking bonds in reactants	Endothermic process
Making bonds in products	Exothermic process

The energy change of reactions (HT only)

AQA Trilogy
C5: Energy Changes

Reaction profiles

Activation energy	Chemical reactions only happen when particles collide with sufficient energy	The minimum amount of energy that colliding particles must have in order to react is called the activation energy.
-------------------	--	--


Overall energy change of a reaction	Exothermic	Energy released making new bonds is greater than the energy taken in breaking existing bonds.
	Endothermic	Energy needed to break existing bonds is greater than the energy released making new bonds.

Bond energy calculation	Calculate the overall energy change for the forward reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$ Bond energies (in kJ/mol): H-H 436, H-N 391, N≡N 945
	Bond breaking: $945 + (3 \times 436) = 945 + 1308 = 2253$ kJ/mol
	Bond making: $6 \times 391 = 2346$ kJ/mol Overall energy change = $2253 - 2346 = -93$ kJ/mol Therefore reaction is exothermic overall.

Endothermic		Products are at a higher energy level than the reactants. As the reactants form products, energy is transferred from the surroundings to the reaction mixture. The temperature of the surroundings decreases because energy is taken in during the reaction.
Exothermic		Products are at a lower energy level than the reactants. When the reactants form products, energy is transferred to the surroundings. The temperature of the surroundings increases because energy is released during the reaction.

Rate of chemical reaction	<i>This can be calculated by measuring the quantity of reactant used or product formed in a given time.</i>	Rate = $\frac{\text{quantity of reactant used}}{\text{time taken}}$ Rate = $\frac{\text{quantity of product formed}}{\text{time taken}}$
---------------------------	---	---

Quantity	Unit
Mass	Grams (g)
Volume	cm ³
Rate of reaction	Grams per cm ³ (g/cm ³) HT: moles per second (mol/s)




Calculating rates of reactions

Rate of reaction

Factors affecting rates

Factors affecting the rate of reaction	
Temperature	<i>The higher the temperature, the quicker the rate of reaction.</i>
Concentration	<i>The higher the concentration, the quicker the rate of reaction.</i>
Surface area	<i>The larger the surface area of a reactant solid, the quicker the rate of reaction.</i>
Pressure (of gases)	<i>When gases react, the higher the pressure upon them, the quicker the rate of reaction.</i>

Collision theory and activation energy




Collision theory	<i>Chemical reactions can only occur when reacting particles collide with each other with sufficient energy.</i>	Increasing the temperature increases the frequency of collisions and makes the collisions more energetic, therefore increasing the rate of reaction.
Activation energy	<i>This is the minimum amount of energy colliding particles in a reaction need in order to react.</i>	Increasing the concentration, pressure (gases) and surface area (solids) of reactions increases the frequency of collisions, therefore increasing the rate of reaction.

AQA Trilogy C6: The Rate and extent of chemical change

Catalysts

Catalyst	A catalyst changes the rate of a chemical reaction but is not used in the reaction.
Enzymes	These are biological catalysts.
How do they work?	Catalysts provide a different reaction pathway where reactants do not require as much energy to react when they collide.



If a catalyst is used in a reaction, it is not shown in the word equation.

Reversible reactions and dynamic equilibrium

Reversible reactions

Reversible reactions	In some chemical reactions, the products can react again to re-form the reactants.
Representing reversible reactions	$A + B \rightleftharpoons C + D$
The direction	The direction of reversible reactions can be changed by changing conditions: $A + B \xrightleftharpoons[\text{cool}]{\text{heat}} C + D$

Energy changes and reversible reactions


If one direction of a reversible reaction is exothermic, the opposite direction is endothermic. The same amount of energy is transferred in each case.

Changing conditions and equilibrium (HT)

Equilibrium

The relative amounts of reactants and products at equilibrium depend on the conditions of the reaction.

Equilibrium in reversible reactions
When a reversible reaction occurs in apparatus which prevents the escape of reactants and products, equilibrium is reached when the forward and reverse reactions occur exactly at the same rate.



Le Chatelier's Principles	States that when a system experiences a disturbance (change in condition), it will respond to restore a new equilibrium state.	
Changing concentration	If the concentration of a reactant is increased, more products will be formed. If the concentration of a product is decreased, more reactants will react.	
Changing temperature	If the temperature of a system at equilibrium is increased: - Exothermic reaction = products decrease - Endothermic reaction = products increase	
Changing pressure (gaseous reactions)	For a gaseous system at equilibrium: - Pressure increase = equilibrium position shifts to side of equation with smaller number of molecules. - Pressure decrease = equilibrium position shifts to side of equation with larger number of molecules.	

Crude oil	<i>A finite resource</i>	Consisting mainly of plankton that was buried in the mud, crude oil is the remains of ancient biomass.
Hydrocarbons	<i>These make up the majority of the compounds in crude oil</i>	Most of these hydrocarbons are called alkanes.
General formula for alkanes	C_nH_{2n+2}	For example: C_2H_6 C_6H_{14}

Alkanes to alkenes	<i>Long chain alkanes are cracked into short chain alkenes.</i>
Alkenes	<i>Alkenes are hydrocarbons with a double bond (some are formed during the cracking process).</i>
Properties of alkenes	<i>Alkenes are more reactive than alkanes and react with bromine water. Bromine water changes from orange to colourless in the presence of alkenes.</i>

Cracking	<i>The breaking down of long chain hydrocarbons into smaller chains</i>	The smaller chains are more useful. Cracking can be done by various methods including catalytic cracking and steam cracking.
Catalytic cracking	<i>The heavy fraction is heated until vaporised</i>	After vaporisation, the vapour is passed over a hot catalyst forming smaller, more useful hydrocarbons.
Steam cracking	<i>The heavy fraction is heated until vaporised</i>	After vaporisation, the vapour is mixed with steam and heated to a very high temperature forming smaller, more useful hydrocarbons.

Crude oil, hydrocarbons and alkanes

Display formula for first four alkanes

$\begin{array}{c} \text{H} \\ | \\ \text{H}-\text{C}-\text{H} \\ | \\ \text{H} \end{array}$

Methane (C_1H_4)

$\begin{array}{c} \text{H} \quad \text{H} \\ | \quad | \\ \text{H}-\text{C}-\text{C}-\text{H} \\ | \quad | \\ \text{H} \quad \text{H} \end{array}$

Ethane (C_2H_6)

$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ | \quad | \quad | \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\ | \quad | \quad | \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$

Propane (C_3H_8)

$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\ | \quad | \quad | \quad | \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{H} \\ | \quad | \quad | \quad | \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \end{array}$

Butane (C_4H_{10})

Carbon compounds as fuels and feedstock

**AQA Trilogy
C7: Organic Chemistry**



Carbon compounds as fuels and feedstock

Cracking and alkenes

Fractional distillation and petrochemicals

Hydrocarbon chains in oil	Hydrocarbon chains in crude oil come in lots of different lengths.
Boiling points	The boiling point of the chain depends on its length. During fractional distillation, they boil and separate at different temperatures due to this.


Combustion	During the complete combustion of hydrocarbons, the carbon and hydrogen in the fuels are oxidised, releasing carbon dioxide, water and energy.
-------------------	--




Alkenes and uses as polymers	<i>Used to produce polymers. They are also used as the starting materials of many other chemicals, such as alcohol, plastics and detergents.</i>
Why do we crack long chains?	<i>Without cracking, many of the long hydrocarbons would be wasted as there is not much demand for these as for the shorter chains.</i>

Boiling point (temperature at which liquid boils)	<i>As the hydrocarbon chain length increases, boiling point increases.</i>
Viscosity (how easily it flows)	<i>As the hydrocarbon chain length increases, viscosity increases.</i>
Flammability (how easily it burns)	<i>As the hydrocarbon chain length increases, flammability decreases.</i>

Fractions	<i>The hydrocarbons in crude oil can be split into fractions</i>	Each fraction contains molecules with a similar number of carbon atoms in them. The process used to do this is called fractional distillation.
Using fractions	<i>Fractions can be processed to produce fuels and feedstock for petrochemical industry</i>	We depend on many of these fuels; petrol, diesel and kerosene. Many useful materials are made by the petrochemical industry; solvents, lubricants and polymers.



Pure substances	<i>A pure substance is a single element or compound, not mixed with any other substance.</i>	Pure substances melt and boil at specific temperatures. Heating graphs can be used to distinguish pure substances from impure.
------------------------	--	--




Melting point of a pure substance



Melting point of an impure substance

Formulation	<i>A formulation is a mixture that has been designed as a useful product.</i>
How are formulations made?	<i>By mixing chemicals that have a particular purpose in careful quantities.</i>
Examples of formulations.	<i>Fuels, cleaning agents, paints, medicines and fertilisers.</i>



Chromatography	<i>Can be used to separate mixtures and help identify substances.</i>	Involves a mobile phase (e.g. water or ethanol) and a stationary phase (e.g. chromatography paper).
R_f Values	<i>The ratio of the distance moved by a compound to the distance moved by solvent.</i>	$R_f = \frac{\text{distance moved by substance}}{\text{distance moved by solvent}}$
Pure substances	<i>The compounds in a mixture separate into different spots.</i>	This depends on the solvent used. A pure substance will produce a single spot in all solvents whereas an impure substance will produce multiple spots.

Pure substances

Purity, formulations and chromatography


Chromatography

Formulations

**AQA Trilogy
C8: Chemical Analysis**

Identification of common gases

Gas	Test	Positive result
Hydrogen	<i>Burning splint</i>	'Pop' sound.
Oxygen	<i>Glowing splint</i>	Re-lights the splint.
Chlorine	<i>Litmus paper (damp)</i>	Bleaches the paper white.
Carbon dioxide	<i>Limewater</i>	Goes cloudy (as a solid calcium carbonate forms).



Gas	Percentage
Nitrogen	~80%
Oxygen	~20%
Argon	0.93%
Carbon dioxide	0.04%

Proportions of gases in the atmosphere	Algae and plants	These produced the oxygen that is now in the atmosphere, through photosynthesis.	carbon dioxide + water → glucose + oxygen $6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2$
Oxygen in the atmosphere		First produced by algae 2.7 billion years ago.	Over the next billion years plants evolved to gradually produce more oxygen. This gradually increased to a level that enabled animals to evolve.

Volcano activity 1 st Billion years	Billions of years ago there was intense volcanic activity	This released gases (mainly CO ₂) that formed to early atmosphere and water vapour that condensed to form the oceans.
Other gases	Released from volcanic eruptions	Nitrogen was also released, gradually building up in the atmosphere. Small proportions of ammonia and methane also produced.
Reducing carbon dioxide in the atmosphere	When the oceans formed, carbon dioxide dissolved into it	This formed carbonate precipitates, forming sediments. This reduced the levels of carbon dioxide in the atmosphere.

The Earth's early atmosphere

How oxygen increased

How carbon dioxide decreased

Composition and evolution of the atmosphere

AQA Trilogy
C9: Chemistry of the atmosphere

Common atmospheric pollutants

CO₂ and methane as greenhouse gases

Greenhouse gases

Carbon dioxide, water vapour and methane	Examples of greenhouse gases that maintain temperatures on Earth in order to support life
The greenhouse effect	Radiation from the Sun enters the Earth's atmosphere and reflects off of the Earth. Some of this radiation is re-radiated back by the atmosphere to the Earth, warming up the global temperature.

Carbon footprints
The total amount of greenhouse gases emitted over the full life cycle of a product/event. This can be reduced by reducing emissions of carbon dioxide and methane.

Global climate change

Human activities and greenhouse gases

Carbon dioxide	Human activities that increase carbon dioxide levels include burning fossil fuels and deforestation.
Methane	Human activities that increase methane levels include raising livestock (for food) and using landfills (the decay of organic matter released methane).
Climate change	There is evidence to suggest that human activities will cause the Earth's atmospheric temperature to increase and cause climate change.

Properties and effects of atmospheric pollutants

Carbon monoxide	Toxic, colourless and odourless gas. Not easily detected, can kill.
Sulfur dioxide and oxides of nitrogen	Cause respiratory problems in humans and acid rain which affects the environment.
Particulates	Cause global dimming and health problems in humans.

Effects of climate change
Rising sea levels
Extreme weather events such as severe storms
Change in amount and distribution of rainfall
Changes to distribution of wildlife species with some becoming extinct

Atmospheric pollutants from fuels

Combustion of fuels	Source of atmospheric pollutants. Most fuels may also contain some sulfur.
Gases from burning fuels	Carbon dioxide, water vapour, carbon monoxide, sulfur dioxide and oxides of nitrogen.
Particulates	Solid particles and unburned hydrocarbons released when burning fuels.



Corrosion	<i>The destruction of materials by chemical reactions with substances in the environment</i>	An example of this is iron rusting; iron reacts with oxygen from the air to form iron oxide (rust) water needs to be present for iron to rust.
Preventing corrosion	<i>Coatings can be added to metals to act as a barrier</i>	Examples of this are greasing, painting and electroplating. Aluminium has an oxide coating that protects the metal from further corrosion.
Sacrificial corrosion	<i>When a more reactive metal is used to coat a less reactive metal</i>	This means that the coating will react with the air and not the underlying metal. An example of this is zinc used to galvanise iron.

Corrosion and its prevention

Alloys are useful materials

Alloys	<i>A mixture of two elements, one of which must be a metal e.g. Bronze is an alloy of copper and tin and Brass is an alloy of copper and zinc.</i>
Gold carats	<i>Gold jewellery is usually an alloy with silver, copper and zinc. The carat of the jewellery is a measure of the amount of gold in it e.g. 18 carat is 75% gold, 24 carat is 100% gold.</i>
Steels	<i>Alloys of iron, carbon and other metals.</i>
	<i>High carbon steel is strong but brittle.</i>
	<i>Low carbon steel is softer and easily shaped.</i>
	<i>Steel containing chromium and nickel (stainless) are hard and corrosion resistant.</i>
	<i>Aluminium alloys are low density.</i>

NPK fertilisers	<i>These contain nitrogen, phosphorous and potassium</i>	Formulations of various salts containing appropriate percentages of the elements.
Fertiliser examples	<i>Potassium chloride, potassium sulfate and phosphate rock are obtained by mining</i>	Phosphate rock needs to be treated with an acid to produce a soluble salt which is then used as a fertiliser. Ammonia can be used to manufacture ammonium salts and nitric acid.

Production and uses of NPK fertilisers

Using materials

**AQA Trilogy
C10: Using Resources**

The Haber process and the use of NPK fertilisers

Ceramics, polymers and composites

Polymers	Thermosetting	polymers that do not melt when they are heated.
	Thermosoftening	polymers that melt when they are heated.

Composite materials	<i>A mixture of materials put together for a specific purpose e.g. strength</i>	Soda-lime glass, made by heating sand, sodium carbonate and limestone.
		Borosilicate glass, made from sand and boron trioxide, melts at higher temperatures than soda-lime glass.
		MDF wood (woodchips, shavings, sawdust and resin)
		Concrete (cement, sand and gravel)
Ceramic materials	<i>Made from clay</i>	Made by shaping wet clay and then heating in a furnace, common examples include pottery and bricks.
Polymers	<i>Many monomers can make polymers</i>	These factors affect the properties of the polymer. Low density (LD) polymers and high density (HD) polymers are produced from ethene. These are formed under different conditions.
The Haber process	<i>Used to manufacture ammonia</i>	Ammonia is used to produce fertilisers Nitrogen + hydrogen \rightleftharpoons ammonia
Raw materials	<i>Nitrogen from the air while hydrogen from natural gas</i>	Both of these gases are purified before being passed over an iron catalyst. This is completed under high temperature (about 450°C) and pressure (about 200 atmospheres).
Catalyst	<i>Iron</i>	The catalyst speeds up both directions of the reaction, therefore not actually increasing the amount of valuable product.

Phosphate rock	
Treatment	Products
Nitric acid	<i>The acid is neutralised with ammonia to produce ammonium phosphate, a NPK fertiliser.</i>
Sulfuric acid	<i>Calcium phosphate and calcium sulfate (a single superphosphate).</i>
Phosphoric acid	<i>Calcium phosphate (a triple superphosphate).</i>

The Haber process – conditions and equilibrium	
Pressure	<i>The reactants side of the equation has more molecules of gas. This means that if pressure is increased, equilibrium shifts towards the production of ammonia (Le Chatelier's principle). The pressure needs to be as high as possible.</i>
Temperature	<i>The forward reaction is exothermic. Decreasing temperature increases ammonia production at equilibrium. The exothermic reaction that occurs releases energy to surrounding, opposing the temperature decreases. Too low though and collisions would be too infrequent to be financially viable.</i>

The Haber process

