



Year 9 AREA OF CIRCLES AND PART CIRCLES





Year 9 PYTHAGORAS AND TRIGONOMETRY

Key Concepts

Pythagoras' theorem and basic trigonometry both only work with right angled triangles.

Pythagoras' Theorem – used to find a missing length when two sides are known $a^2 + b^2 = c^2$ c is always the hypotenuse (longest side)





A hegartymaths

498-499,

509-515



m00.8 (6:283W2NA b) 2.94m c) 55.15° d) 2.34cm





VOLUME AND SURFACE AREAS OF CYLINDERS

Year 9

Key Concepts

A **cylinder** is a **prism** with the cross section of a circle.



The **volume** of a cylinder is calculated by $\pi r^2 h$ and is the space inside the 3D shape

The **surface area** of a cylinder is calculated by $2\pi r^2 + \pi dh$ and is the total of the areas of all the faces on the shape.

A hegartymaths

572, 586

a) **Volume** $V = \pi \times r^2 \times h$

10cm

From the diagram calculate:

4cm

 $V = \pi \times 4^2 \times 10$

Key Words

Cylinder Surface Area

Radius

Diameter Pi Volume

Prism

 $V = 160\pi$ Or = 502.65 cm^3

Examples

b) Surface Area – You can use the net of the shape to help you

Area of two circles = $2 \times \pi \times r^2$ = $2 \times \pi \times 4^2$ = 32π

Area of rectangle $= \pi \times d \times h$ $= \pi \times 8 \times 10$ $= 80\pi$



Surface Area = $32\pi + 80\pi$ = 112π or = $351.86cm^3$

Calculate the volume and surface area of this cylinder



 $^{\circ}$ mo10.768 or $\pi 305$ = $\pi 305$ Surface area = 308π or $\pi 357$ = $\pi 308$ SNA



from rounding.

decimal place:

Year 9 BOUNDARIES

X

 $UB_1 \times UB_2$

 $LB_1 \times LB_2$

Examples

When completing calculations involving boundaries we are aiming to find the greatest or smallest answer.

A restaurant provides a cuboid stick of butter to each table. The dimensions are 30mm by 30mm by 80mm, correct to the nearest 5mm. Calculate the upper and lower bounds of the volume of the butter.

+

 $UB_1 + UB_2$

 $LB_1 + LB_2$

 $UB_1 - LB_2$

 $LB_1 - UB_2$

 $Volume = l \times w \times h$

Upper bound

answer

Lower bound

answer

 $Upper \ bound = 32.5 \times 82.5 \times 32.5 \\ = 87140.63mm^3 \\ Lower \ bound = 27.5 \times 77.5 \times 27.5 \\ = 58609.38mm^3$

 $D = \frac{x}{v}$

÷

 $UB_1 \div LB_2$

 $LB_1 \div UB_2$

x = 99.7 correct to 1 decimal place.y = 67 correct to 2 significant figures.Work out an upper and lower bounds for *D*.

Upper bound
$$D = \frac{99.75}{66.5} = 1.5$$

Lower bound
$$D = \frac{99.65}{67.5} = 1.48$$

Jada has 100 litres of oil, correct to the nearest litre.
The oil is poured into tins of volume 1.5 litres, correct to one decimal place.
Calculate the upper and lower bounds for the number of tins that can be filled.

2) There are 110 identical marbles in a bag. A marble is taken and weighed as 15.6 g to the nearest tenth of a gram. Find the upper and lower bounds for the weight of all the marbles.

ANSWERS: 1) LB = 69.3 \approx 69 UB = 64.2 \approx 64 2) LB = 1721.5 g UB = 1721.5 g

significant figures: $355 \le x < 365$ E.g. State the boundaries of 4.5 when it has been rounded to 2

Key Concepts

The boundaries of a number derive

E.g. State the boundaries of 360

when it has been rounded to 2

 $4.45 \le x < 4.55$

These boundaries can also be called the **error interval** of a number.

A hegartymaths

137-139,

411

Key Words Bound Upper Lower Accuracy Rounding